

Report Guidelines

Title: Developing seasonal streamflow forecasts to inform surface water management in Oklahoma

Authors' Names and Affiliations:

Tyson Ochsner, Professor, Plant and Soil Sciences, Oklahoma State University, tyson.ochsner@okstate.edu, 405-744-3627

Briana Wyatt, Doctoral Candidate, Department of Plant and Soil Sciences, 371 Stillwater, OK. Email: briana.wyatt@okstate.edu Phone: NA

Erik Krueger, Research Associate, Plant and Soil Sciences, Oklahoma State University, erik.krueger@okstate.edu, 405-744-3627

Eric Jones, Development and Operations Hydrologist, Arkansas-Red Basin River Forecast Center, National Weather Service, eric.jones@noaa.gov, 918-832-4109

Start Date: 06/16/2018

End Date: 06/17/2019

Congressional District: Oklahoma District 3

Focus Category: HYDROL, M&P, MOD, SW, WQN

Descriptors: soil moisture, streamflow forecasting

Students:

Student Status	Number	Disciplines
Undergraduate		
M.S.		
Ph.D.	1	Soil Science
Post Doc		
Total		

Principal Investigators:

Tyson Ochsner

Professor, Department of Plant and Soil Sciences, 371 Stillwater, OK.
Email: tyson.ochsner@okstate.edu

Briana Wyatt

Doctoral Candidate, Department of Plant and Soil Sciences, 371 Stillwater, OK.

Email: briana.wyatt@okstate.edu

Erik Krueger

Research Associate, Department of Plant and Soil Sciences, 371 Stillwater, OK.

Email: erik.krueger@okstate.edu

Eric Jones

Operations Hydrologist, NWS Arkansas/Red River Forecast Center Tulsa, OK.

Email: eric.jones@noaa.gov

Publications and Presentations:

- Wyatt, B.M., T.E. Ochsner, and E.S. Krueger. 2019. Improving seasonal streamflow forecasts by incorporating soil moisture data. SSSA International Soils Meeting. San Diego, CA.
- Wyatt, B.M., T.E. Ochsner, E.S. Kreuger, and E.T. Jones. 2019. Improving seasonal streamflow forecasts by incorporating soil moisture data. National Soil Moisture Network annual meeting. Manhattan, KS.
- Wyatt, B.M., T.E. Ochsner, E.S. Kreuger, and E.T. Jones. 2019. Improving seasonal streamflow forecasts by incorporating soil moisture data. Oklahoma State University Plant and Soil Sciences Department Research Symposium. Stillwater, OK.
- Wyatt, B.M., T.E. Ochsner, E.S. Kreuger, and E.T. Jones. 2019. Improving seasonal streamflow forecasts by incorporating soil moisture data. Oklahoma Clean Lakes and Watersheds Association Annual Meeting. Stillwater, OK.
- Wyatt, B.M., T.E. Ochsner, E.S. Kreuger, and E.T. Jones. 2019. Improving seasonal streamflow forecasts by incorporating soil moisture data. Oklahoma State University Plant and Soil Sciences Department Seminar. Stillwater, OK.
- Wyatt, B.M., T.E. Ochsner, E.S. Krueger, and E. Jones. 2018. Improving seasonal streamflow forecasts by incorporating soil moisture data. National Institutes for Water Resources Regional Symposium. Lincoln, NE.
- Wyatt, B.M., T.E. Ochsner, E.S. Krueger, and E. Jones. 2018. Improving seasonal streamflow forecasts by incorporating soil moisture data. Marena, Oklahoma In-Situ Sensor Testbed (MOISST) annual meeting. Lincoln, NE.
- Wyatt, B.M., T.E. Ochsner, E.S. Krueger, and E. Jones. 2018. Improving seasonal streamflow forecasts to inform surface water management in Oklahoma by incorporating soil moisture data. Oklahoma State University Plant and Soil Sciences Department Research Symposium. Stillwater, OK.

Problem and Research Objectives:

Worldwide, an increasingly variable climate and the growing water demands of a rising world population continually threaten the security and sustainability of surface water resources essential for agricultural production, domestic and industrial water use, recreation, and other beneficial uses (Brekke et al., 2010; Wood, 2007; Garbrecht et al., 2004). These threats are amplified by the relatively low skill of many hydrological forecasting systems for adequately predicting future streamflows (Nash and Sutcliffe, 1970). While operational models applied in snow-dominated watersheds have been widely studied and improved (Pagano et al., 2004), seasonal forecasting methods (i.e., forecasts which estimate streamflow volumes over several months) for rainfall-dominated watersheds remain less skillful, reducing their efficacy for water resource management (Cuo et al., 2011; Pagano et al., 2004). Increasing the accuracy of seasonal streamflow forecasts in rainfall-dominated watersheds is critical for improving reservoir operations, drought management, sustainable water use, hydropower production, and irrigated agriculture (Raff et al., 2013; Maurer and Lettenmaier, 2004).

An alternative forecasting method commonly used to predict streamflow in the snow-dominated, mountainous Western U.S. is principal components analysis and regression (or principle components regression, hereafter referred to as PCR), a rigorous statistical method utilized by the Natural Resources Conservation Service (NRCS) (Garen, 1992; Garen and Pagano, 2007). Harpold et al. (2017) showed that the inclusion of soil moisture data in PCR analysis improved operational streamflow forecast skill in 12 snow-dominated watersheds in the Western U.S. Our objective was to evaluate the potential improvements from including in situ soil moisture data in PCR-based streamflow forecasts in *rainfall-dominated* watersheds.

Methodology:

Our study focused on two Oklahoma watersheds- Fort Cobb and Little Washita- as well as two watersheds outside the state- Walnut Gulch, Arizona and Little River, Georgia (Figure 1). Similar to Harpold et al. (2017), we chose to use a two-step PCR analysis in order to distinguish the presumably first-order control of antecedent precipitation on streamflow from the presumably second-order control of soil moisture. In our case, this involved an initial PCR analysis using only antecedent precipitation data as inputs (hereafter called the “baseline” scenario) to estimate seasonal streamflow totals and a secondary PCR (hereafter called the “two-step” scenario) which estimated the residuals between estimated and observed streamflow volumes from the baseline scenario using five soil moisture metrics- volumetric water content, percent saturation, total storage, available storage, and the Soil Moisture Index (SMI). Baseline and two-step models were each used to predict streamflow totals during the four months of greatest streamflow in each watershed at lead times of up to 3 months.




Figure 1. The Walnut Gulch (a) watershed in Arizona, Fort Cobb (b) and Little Washita (c) watersheds in Oklahoma, and the Little River watershed in Georgia (d). Yellow stars represent the locations of co-located precipitation and soil moisture monitoring stations, except for in the Walnut Gulch watershed, where yellow stars indicate the location of soil moisture monitoring stations and black circles indicate the location of precipitation stations. Streamflow gauges are indicated by black triangles in all watersheds, except at Fort Cobb where lake inflow levels were used rather than stream gauge data and where the Fort Cobb Dam is marked by a black bar.

Principal Findings and Significance:

Baseline forecasts made using only antecedent precipitation data were only able to produce forecasts at the 0-month lead time in two of the four watersheds, and no forecasts could be made at longer lead times. The two 0-month baseline forecasts that were made were for the Fort Cobb and Little River watersheds, and explained 27% and 19% of seasonal streamflow variability, respectively. Additionally, both forecasts were classified as unsatisfactory based on performance criteria described by Moriasi et al. (2007).

Conversely, the inclusion of soil moisture data in the two-step forecasts led to forecasts being made in all watersheds at all lead times. These forecasts explained between 35% to 87% of seasonal streamflow variability, with 0-month forecasts explaining an average of 78% of variability. Of forecasts made using soil moisture data, 88% were rated as satisfactory or better based on performance criteria described by Moriasi et al. (2007). Forecasts for all watersheds are shown in Figure 2, where each point represents one water year's streamflow volumes. Figure 2 demonstrates how the inclusion of soil moisture data in PCR forecasts improves forecast accuracy, particularly in years when baseline forecast error is high. These results represent the first evidence that the PCR method can produce accurate seasonal streamflow forecasts in rainfall-dominated regions and that including soil moisture data in the PCR model increases forecast accuracy over forecasts made using antecedent precipitation data alone.




Figure 2. Yearly observed and predicted seasonal streamflow volumes for baseline (black squares) and two-step (triangles) forecasts in the Walnut Gulch (a), Fort Cobb (b), Little Washita (c), and Little River (d) watersheds at the 0-month lead time. Dashed line is 1:1 line. Baseline forecasts were based only on cumulative precipitation prior to the forecast date (ΣP), whereas two-step forecasts included both ΣP and soil moisture (SM) information. Baseline forecasts for the Walnut Gulch and Little River watersheds did not meet the criteria for statistical validity, but baseline streamflow predictions are shown here to demonstrate improvements due to soil moisture data inclusion.

References

- Brekke, L. D., D. Garen, K. Werner, and D. Laurine. 2010. Projecting climate change impacts on seasonal water supply forecasting error. Preprints, 18th Conf. on Applied Climatology/22nd Conf. on Climate Variability and Change/ 24th Conf. on Hydrology, Atlanta, GA-LR, Amer. Meteor. Soc., J15.4. [Available online at https://ams.confex.com/ams/90annual/techprogram/paper_162386.htm.]
- Moriasi, D., et al. 2007. Model evaluation guidelines for systematic quantification of accuracy in watershed simulations. *Trans. Asabe* 50(3): 885-900.

- Harpold, A. A., et al. 2017. Does Including Soil Moisture Observations Improve Operational Streamflow Forecasts in Snow-Dominated Watersheds? JAWRA Journal of the American Water Resources Association 53(1): 179-196.
- Garen, D. C. 1992. Improved techniques in regression-based streamflow volume forecasting. Journal of Water Resources Planning and Management 118(6): 654-670.
- Garen, D.C., and T. C. Pagano, 2007: Statistical techniques used in the VIPER water supply forecasting software. NRCS-USDA Engineering-Snow Survey and Water Supply Forecasting Tech. Note 210-2, 18 pp. [Available online at http://www.wcc.nrcs.usda.gov/ftpref/downloads/factpub/wsf/technotes/Tech_note_statistical_techniques_in_Viper.pdf.]
- Raff, D., L. Brekke, K. Werner, A.W. Wood and K. White. 2013. Short-term water management decisions: User needs for improved climate, weather, and hydrologic information. In: U.S. Army Corps of Engineers, Bureau of Reclamation and National Oceanic and Atmospheric Administration, editors, National Technical Information Service, Springfield, Virginia. p. 233.
- Maurer, E. P. and D. P. Lettenmaier. 2004. Potential Effects of Long-Lead Hydrologic Predictability on Missouri River Main-Stem Reservoirs. Journal of Climate 17(1): 174-186.
- Pagano, T., D. Garen, and S. Sorooshian. 2004. Evaluation of Official Western U.S. Seasonal Water Supply Outlooks, 1922–2002. Journal of Hydrometeorology 5(5):896-909.
- Cuo, L., T.C. Pagano, and Q.J. Wang. 2011. A Review of Quantitative Precipitation Forecasts and Their Use in Short- to Medium-Range Streamflow Forecasting. Journal of Hydrometeorology 12:713-728.
- Nash, J. E. and J. V. Sutcliffe. 1970. River flow forecasting through conceptual models part I — A discussion of principals. Journal of hydrology 10(3): 282-290.
- Garbrecht, J., et al. 2004. Trends in Precipitation, Streamflow, and Evapotranspiration in the Great Plains of the United States. Journal of Hydrologic Engineering 9(5): 360-367.
- Wood, A. W. 2007. The effects of climate change on water supply forecasting in the Feather River basin. Preprints, Fourth Annual California Climate Change Conf., Sacramento, CA, California Energy Commission.